

**U.S. PLANT PATENT APPLICATION OF**

**HEINRICH WESTHOFF**

**FOR: LOBELIA PLANT NAMED**

**‘WESLOWHITE’**

WESTHOFF, Heinrich

APPLICANT: HEINRICH WESTHOFF

TITLE: LOBELIA PLANT NAMED 'WESLOWHITE'

BOTANICAL CLASSIFICATION/CULTIVAR DESIGNATION:

*Lobelia X hybrida* cultivar WESLOWHITE

5

## BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cultivar of Lobelia plant, botanically known as *Lobelia erinus*, and hereinafter referred to by the name 'Weslowwhite'.

10

The new Lobelia is a product of a planned breeding program conducted by the Inventor in Südlohn, Germany. The objective of the breeding program was to develop new Lobelia cultivars with continuous flowering, white flower color and tolerance to high temperatures.

15

The new Lobelia originated from a cross-pollination made by the Inventor in 2002 of a proprietary selection of *Lobelia erinus* identified as code number 01P050, not patented, as the female, or seed, parent with the *Lobelia erinus* cultivar Weslobigblue, disclosed in U.S. Plant Patent number 12,634, as the male, or pollen, parent. The new Lobelia was discovered and selected by the Inventor from within the resultant

progeny from the above-mentioned cross-pollination in a controlled environment in Südlohn, Germany in 2002.

Asexual reproduction since 2002 of the new cultivar by terminal cuttings in a controlled environment in Südlohn, Germany, has shown  
5 that the unique features of this new Lobelia are stable and reproduced true to type in successive generations.

#### SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Weslowwhite'. These  
10 characteristics in combination distinguish 'Weslowwhite' as a new and distinct cultivar:

1. Cascading plant habit.
2. Freely branching habit with short internodes; dense and bushy plant form.
- 15 3. Continuously and freely flowering habit.
4. Densely pubescent leaves.
5. White- colored flowers.
6. Relatively tolerant to high temperatures.

Plants of the new Lobelia have larger flowers than plants of the  
20 female parent, a proprietary selection identified as code number 01P050.

In addition, plants of the new Lobelia produce flowers that have colored markings on the petals whereas plants of the female parent selection do not have colored markings on the petals. Plants of the new Lobelia have foliage with dense pubescence compared to the smooth foliage of plants of the male parent, the cultivar Weslobigblue. In addition, plants of the new Lobelia differ from plants of the cultivar Weslobigblue in leaf shape and flower color.

Plants of the new Lobelia differ primarily from plants of the cultivar Weslolav, disclosed in a U.S. Plant Patent application filed concurrently, primarily in flower color.

Plants of the cultivar Weslowhite can be compared to the cultivar Weslowei, disclosed in U.S. Plant Patent number 12,708. However in side-by-side comparisons conducted in Südlohn, Germany, plants of the cultivar Weslowhite and the cultivar Weslowei differed in the following characteristics:

1. Leaves and stems of plants of the new Lobelia were more pubescent than plants of the cultivar Weslowei.
2. Plants of the new Lobelia flowered more continuously than plants of the cultivar Weslowei.

3. Flowers of the new Lobelia were shaped differently than flowers of the cultivar Weslowei.
4. Plants of the new Lobelia had flowers with different color markings than flowers of plants of the cultivar Weslowei.
5. Plants of the new Lobelia were more tolerant to high temperatures than plants of the cultivar Weslowei.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description, which accurately describe the actual colors of the new Lobelia. The photograph at the top of the sheet comprises a close-up view of a typical individual flower of 'Weslowwhite'. The photograph at the bottom of the sheet comprises side view of a typical plant of 'Weslowwhite' grown in a hanging basket container.

#### DETAILED BOTANICAL DESCRIPTION

Plants of the cultivar Weslowwhite have not been observed under all possible environmental conditions. The phenotype may vary

somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

In the following description, color references are made to the Royal Horticultural Society Colour Chart, 2001 Edition, except where  
5 general terms of ordinary dictionary significance are used. Plants used for the description were grown in a glass-covered greenhouse and conditions that closely approximate commercial production conditions during the spring and summer in Südlöhn, Germany. Plants used for the above-mentioned photographs and following description were grown as  
10 one plant per 12-cm container or three plants per 25-cm hanging basket container. During the production of the plants, day temperatures ranged from 20 to 25°C and night temperatures ranged from 16 to 18°C. Plants were pinched once during the production period by removing about 1 to 2 cm of the uppermost apical growing tip. Plants were about 20 weeks  
15 from planting when the photographs and description were taken.

#### BOTANICAL CLASSIFICATION:

*Lobelia erinus* cultivar Weslowhite.

#### PARENTAGE:

Female parent: Proprietary selection of *Lobelia erinus* identified  
20 as code number 01P050, not patented.

Male parent: *Lobelia erinus* cultivar Weslobigblue, disclosed in  
U.S. Plant Patent number 12,634.

PROPAGATION:

Type cutting: Terminal vegetative cuttings.

5 Time to initiate roots: About 18 to 21 days at 20°C.

Time to develop roots: About 20 to 28 days at 20°C.

Root description: Fine, fibrous and well-branched.

PLANT DESCRIPTION:

10 Plant form/habit: Cascading flowering plants with white- colored  
flowers. Lateral shoots outwardly spreading; plants uniform with  
dense foliage. Foliage and stems are very pubescent. Freely  
branching with lateral branches forming at every node; dense and  
bushy plant habit. Pinching plants will enhance branching.  
Moderately vigorous growth habit.

15 Usage: Appropriate for hanging baskets, window boxes and patio  
containers.

Plant height (soil level to top of plant plane): About 18.5 cm.

Plant length (soil level to lateral branches apices): About 70 cm.

Plant diameter: About 35 to 40 cm.

Lateral branch description:

Length: About 17 cm.

Diameter: About 1.7 mm.

Internode length: About 1.5 cm.

5 Texture: Densely pubescent.

Color: 146A and 146B.

Foliage description:

Arrangement: Alternate; simple.

Basal leaves:

10 Length: About 4.6 cm.

Width: About 3.2 cm.

Shape: Ovate.

Apex: Rounded.

Base: Attenuate.

15 Margin: Crenate.

Petiole length: About 1.6 mm.

Mid-plant leaves:

Length: About 3.6 cm.

Width: About 2.3 cm.

20 Shape: Ovate.



Apex: Rounded.

Base: Attenuate.

Margin: Nearly entire.

Petiole length: About 9.5 mm.

5

Apical leaves:

Length: About 2.6 cm.

Width: About 5 mm.

Shape: Oblanceolate to linear.

Apex: Acute.

10

Base: Attenuate.

Margin: Entire.

Petiole length: Petioles not observed.

Texture, all leaves, upper and lower surfaces: Densely pubescent.

15

Color, all leaves:

Developing foliage, upper surface: 147A.

Developing foliage, lower surface: 147B.

Fully developed foliage, upper surface: 147A;  
venation, 147B.

Fully developed foliage, lower surface: 147B to  
147C; venation, 147B.

**FLOWER DESCRIPTION:**

5 Flower type and habit: Flowers arranged singly at lateral apices.  
Flowers held mostly outwardly. Flowers persistent. Older  
flowers are overgrown by new flowers and foliage. Freely and  
continuously flowering. Flowers not fragrant.

Flower shape: Tubular with three larger lower petals and two  
upright petals.

10 Natural flowering season: Spring until frost in the autumn.

Flower longevity on the plant: Longevity of individual flowers  
is highly dependent on weather conditions; typically three to ten  
days.

Flower size:

15 Diameter: About 1.8 cm.

Depth (height): About 2.2 cm.

Tube length: About 1 cm.

Throat diameter, distal end: About 4.7 mm.

Tube diameter, proximal end: About 2.1 mm.

Flower buds:

Length: About 1.45 cm.

Diameter: About 2.6 mm.

Shape: Oblong.

5           Color: Base and mid-section, 145D; towards the apex,  
145C.

Petals:

Arrangement: Single whorl of five petals, fused; three  
larger lower petals and two smaller upper petals.

10           Three lower petals:

Shape: Obovate.

Length, above throat: About 1.3 cm.

Width: About 5.5 mm.

Two upper petals:

15           Shape: Spatulate.

Length, above throat: About 7 mm.

Width: About 2 mm.

Upper and lower petals:

Apex: Cuspidate.

20           Margin: Entire.

Texture, upper and lower surfaces: Smooth, satiny.

Color:

When opening, upper and lower surfaces:

155B.

5 Fully opened, upper and lower surfaces:

155C.

Throat: 69C to 69D; stripes, 144B, and spots,  
76A.

Tube: 69C to 69D; stripes, 144A.

10 Sepals:

Arrangement: Single whorl of five sepals, star-shaped  
calyx.

Length: About 8 mm.

Width: About 1 mm.

15 Shape: Elliptic.

Apex: Acute.

Margin: Entire.

Texture, upper and lower surfaces: Densely pubescent.

Color, upper and lower surfaces: 147A.

Peduncles:

Appearance: Wiry, flexible.

Length: About 2.8 cm.

Diameter: About 6 mm.

5 Texture: Densely pubescent.

Color: 137A.

Reproductive organs:

Stamens:

Quantity per flower: About five, fused.

10 Anther length: About 2.6 mm.

Anther diameter: About 1.8 mm.

Anther texture: Pubescent.

Anther color: 79A to 84D.

Pollen amount: Moderate.

15 Pollen color: 8A.

Pistils:

Quantity per flower: One.

Pistil length: About 1.1 cm.

Stigma shape: Two- parted, ovate.

20 Stigma texture: Pubescent.

Stigma color, immature: 77B.

Stigma color, mature: 77B.

Style length: About 6 mm.

Style color: 144A.

5 Ovary color: 137B and 137C.

Seed/fruit: Seed and fruit production have not been observed.

#### DISEASE/PEST RESISTANCE:

10 Plants of the new Lobelia have not been noted to be resistant to pathogens and pests common to Lobelia.

#### TEMPERATURE TOLERANCE:

Plants of the new Lobelia have been observed to tolerate temperatures ranging from 2 to 30°C.